



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

275. Proposed by PROFESSOR WILLIAM HOOVER, Ph. D., Athens, Ohio.

An hyperbola is drawn touching the axes of an ellipse, and the asymptotes of the hyperbola touch the ellipse. Prove that the center of the hyperbola lies on one of the equal conjugate diameters of the ellipse.

276. Proposed by G. I. HOPKINS, Manchester, N. H.

ABC is an equilateral triangle whose vertices are the centers of circles with radius AB , and H is the center of the arc AB . From F , the point of intersection of the circles whose centers are A and C , a line is drawn through H to the circumference CAN . Draw BN , and prove that the angle ABN is an angle of a regular pentagon.

MECHANICS.

186. Proposed by R. D. CARMICHAEL, Hartselle, Alabama.

A point P keeps at uniform distance from and moves with uniform angular velocity around a point Q which is in harmonic motion, making one revolution while Q swings to and fro. If P is in the line of the path of Q and on the same side of the center of that path with Q when Q is at the extremity of the path, what is the locus of P ?

UNSOLVED PROBLEMS.

NOTE. The following problems still remain unsolved (in our columns).

Algebra, 179. Proposed by DR. L. E. DICKSON, The University of Chicago.

Find the roots of the algebraically solvable quintic equation

$$x^5 + qx^2 + px + \frac{1}{5} \left[\frac{q^2}{p} - \frac{p^3}{5q} \right] = 0.$$

Geometry, 267. Proposed by W. W. LANDIS, Dickinson College, Carlisle, Pa.

Prove that every orthogonal system of circles is an isothermal system.

Group Theory, 9. Proposed by DR. L. E. DICKSON, The University of Chicago.

Does there exist a triply transitive group on m letters of order $m(m-1)(m-2)$ other than the linear fractional group in the Galois Field of order $p^n = m-1$ and the group 720_3 on ten letters (Cole, *Quarterly Journal*, 1895, p. 44)? This question relates to Problem 99, MONTHLY, March, 1900.

Miscellaneous, 151. Proposed by W. J. GREENSTREET, M. A., Stroud, England.

$$\text{Sum the series } \sum_{r=1}^{r=m} \operatorname{cosec} \left[\frac{2r-1}{4m} \pi + \theta \right] \operatorname{cosec} \left[\frac{2r-1}{4m} \pi - \theta \right].$$